

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 June 2002 (27.06.2002)

PCT

(10) International Publication Number
WO 02/49616 A1

(51) International Patent Classification⁷: **A61K 9/12, 47/26**

(21) International Application Number: **PCT/SE01/02853**

(22) International Filing Date:
19 December 2001 (19.12.2001)

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
(XX)4750-6 19 December 2000 (19.12.2000) **SE**

GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

(71) Applicant (for all designated States except US): **ASTRAZENECA AB [SE/SE]; S-151 85 Södertälje (SE).**

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW**, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BUCKTON, Graham [GB/GB];** University of London, School of Pharmacy, 29-39 Brunswick Square, London WC1N 1AZ (GB). **COLUMBANO, Angela [GB/GB];** AstraZeneca R & D Charnwood, Bakewell Road, Loughborough, Leics LE11 5RH (GB). **GROSVENOR, Martin [GB/GB];** AstraZeneca R & D Charnwood, Bakewell Road, Loughborough, Leics LE11 5RH (GB). **WIKLEY, Philip [GB/GB];** Fine Agrochemicals, Hill End House, Whittington, Worcestershire WR5 2RQ (GB).

(74) Agent: **GLOBAL INTELLECTUAL PROPERTY; AstraZeneca AB, S-151 85 Södertälje (SE).**

Published:

— with international search report

(81) Designated States (national): **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,**

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **PHARMACEUTICAL AEROSOL FORMULATION**

(57) Abstract: The invention relates to a pharmaceutical aerosol formulation comprising a surfactant that is an alkyl-polyglycoside of formula I, wherein DP is the average degree of polymerisation and has a value of from 1 to 4, and R is an alkyl chain or a mixture of alkyl chains having a chain length of from 6 to 22 carbon atoms; or a derivative thereof for the administration of a medicament for inhalation.

WO 02/49616 A1

Pharmaceutical aerosol formulation

The present invention relates to a pharmaceutical aerosol formulation for the administration of a medicament for inhalation.

5

Drugs for treating respiratory and nasal disorders are frequently administered in aerosol formulations through the mouth or nose. One widely used method for dispensing such an aerosol formulation involves making a suspension formulation of the drug as a finely divided powder in a liquefied propellant gas. Alternatively a solution formulation can be prepared where the drug is dissolved in a propellant system, perhaps containing solubilisers and co-solvents to aid dissolution of the drug. Pressurised metered dose inhalers (pMDIs) are normally used to dispense such formulations to a patient. Surface active agents, or surfactants, are commonly included in order to aid dispersion of the drug in the propellant and to prevent aggregation of the drug particles, and to improve lubrication. In solution formulations they are used to help solubilise the drug. Following the implication of chlorofluorocarbon (CFC) propellants in the destruction of the ozone layer, CFCs are being replaced by hydrofluoroalkane (HFA) propellants. However the range of surfactants commonly used in the CFC systems have proven to be generally unsuitable for use in HFA systems. Various alternative surfactants have been proposed.

20


For example, WO92/00061 discloses polyethoxylated surfactants for use with HFA propellants. WO91/11173 discloses fluorinated surfactants. WO91/14422 discloses perfluorinated carboxylic acid surfactants for use with HFA propellants. WO92/00107 discloses the use of a propellant-soluble surfactant with propellant 134a, 1,1,1,2-tetrafluoroethane. WO96/19198 discloses aerosol formulations comprising a HFA propellant, a medicament for inhalation and a surfactant which is a C₈₋₁₆ fatty acid or salt thereof, a bile salt, a phospholipid, or an alkyl saccharide.

25

We have found a class of surfactants which are particularly suitable for use in hydrofluoroalkane propellant systems. Said surfactants are highly surface active and allow the generation of dispersion or solution agree leave solutions in formulations with HFA propellants. Further advantages of the said surfactants are their low ecotoxicity, and the ability to synthesise ranges of surfactants with a range of properties such as Molecular Weight, degree of polymerisation.

35

The present invention provides a pharmaceutical aerosol formulation comprising a hydrofluoroalkane propellant, a medicament for inhalation and a surfactant, characterised in that the surfactant is an alkyl-polyglycoside of formula I:



wherein DP is the average degree of polymerisation and has a value of from 1 to 4, and R is an alkyl chain or a mixture of alkyl chains having a chain length of from 6 to 22 carbon atoms; or a derivative thereof.

- The alkyl chain R may be linear or branched and may, for example, be derived from a primary, secondary or tertiary alcohol.

Derivatives of the alkylpolyglycosides of formula I include butyl ethers, carbonates and sulfo-succinates.

- The nomenclature for alkylpolyglycosides is C_nG_m for "pure surfactants" where n is defined as the number of carbon atoms in the alkyl chain and m the number of glucose units attached as the head group. The term "pure" is a relative one as the pure surfactants are not pure in surface chemistry terms but are more homogeneous than technical grade surfactants. Technical grade surfactants are described by a range of alkyl chain lengths (C_{a-b}) as they are derived from natural alcohols, which have a range of alkyl chain lengths and an average number of glucose groups per molecule which is expressed as a degree of polymerisation

(d.p.). Thus C₁₀₋₁₂ d.p. 1.4 has a range of alkyl chain from decyl to dodecyl and an average of 1.4 glucose units per alkyl chain in the sample.

Alkylpolyglycosides are commercially available. Examples of commercial
5 alkylpolyglycosides suitable for use in the present invention include:-

"Berol AG6202™" from Akzo Nobel, an alkylpolyglycoside of formula I in which R is 2-ethyl-1-hexylglycoside and DP is 1.6;

10 "Glucopon 215CS™" from Henkel, an alkylpolyglycoside of formula I in which R is a mixture of C₈ and C₁₀ alkyl chains in a ratio of 60 C₈ : 40 C₁₀ and DP is 1.5;

"Montanov 68™" from SEPPIC, an alkylpolyglycoside of formula I in which R is a mixture
of C₁₆ and C₈ alkyl chains and DP is 1.2 - 1.3; and

15 "Montanov 202™", from SEPPIC an alkylpolyglycoside of formula I in which R is a mixture of C₂₀ and C₂₂ alkyl chains and DP is 1.2 - 1.3.

n-dodecyl β-D-maltoside (C₁₂G₂) is available from Sigma.

20 C₁₀ d.p 2.7 is available from Akzo-Nobel

C₁₀₋₁₂ d.p. 1.4 and C₁₂₋₁₄ d.p. 1.4 are available from Henkel

25 The propellant may be any of hydrofluoroalkane propellant; preferred propellants include 1,1,1,2-tetrafluoroethane (HFA-134a), 1,1,1,2,3,3,3-heptafluoropropane (HFA-227ea) and a mixture of HFA-134a and HFA-227ea, for example a density matched mixture of HFA-134a and HFA-227ea.

30 Medicaments suitable for inclusion in the formulation of the present invention are any which may be delivered by inhalation.

Suitable inhalable medicaments may include for example β₂-adrenoreceptor agonists for example salbutamol, terbutaline, rimiterol, fenoterol, reproterol, adrenaline, pirbuterol,
35 isoprenaline, orciprenaline, bitolterol, salmeterol, formoterol, clenbuterol, procaterol, broxaterol, picumeterol, TA-2005 and malbuterol; anticholinergic bronchodilators for

example ipratropium bromide, oxitropium and its salts and tiotropium and its salts; glucocorticosteroids for example beclomethasone, fluticasone, budesonide, tipredane, dexamethasone, betamethasone, fluocinolone, triamcinolone acetonide, flunisolide, mometasone and 16, 17-acetals of pregnane derivatives, for example rofleponide palmitate and ciclesonide; anti-allergic medicaments for example sodium cromoglycate and nedocromil sodium; leukotriene antagonists for example, zafirlukast, montelukast, pranlukast, zileuton antihistamines for example terfenadine, cetirizine, loratadine and azelastine; antibiotics, ; pain control substances, for example morphine, codeine, pethidine, etc

Specific compounds that can be used in the formulations of the invention are:

4-Hydroxy-7-[2-[2-[3-(2-phenylethoxy)propylsulphonyl]ethylamino]ethyl]-1,3-benzothiazol-2(3H)-one;

4-Hydroxy-7-[2-[2-[3-(2-phenylethoxy)propoxy]ethylamino]ethyl]-1,3-benzothiazol-2(3H)-one;

N-[2-[2-(4-Hydroxy-2-oxo-3H-1,3-benzothiazol-7-yl)ethylamino]ethyl]-2-(phenylethoxy)ethanesulphonamide;

4-Hydroxy-7-[2-[3-[2-[2-(1-naphthalenyl)ethoxy]ethylsulphonyl]propylamino]ethyl]-1,3-benzothiazol-2(3H)-one; and

3-[2-(4-Hydroxy-2-oxo-3H-1,3-benzothiazol-7-yl)ethylamino]-N-[2-[2-(4-methylphenyl)ethoxy]ethyl]propanesulphonamide,

and pharmaceutically acceptable salts thereof, in particular hydrochloride salts.

Preferred medicaments that can be used in the formulations of the invention include

formoterol, terbutaline, budesonide, a formoterol/budesonide combination (e.g Symbicort®), 4-Hydroxy-7-[2-[2-[3-(2-phenylethoxy)propoxy]ethylamino]ethyl]-1,3-benzothiazol-2(3H)-one hydrochloride and 3-[2-(4-Hydroxy-2-oxo-3H-1,3-benzothiazol-7-yl)ethylamino]-N-[2-[2-(4-methylphenyl)ethoxy]ethyl]propanesulphonamide.

Combinations of medicaments may also be employed, for example formoterol/budesonide; formoterol/fluticasone; formoterol/mometasone; salmeterol/fluticasone; formoterol/tiotropium salts; zafirlukast/formoterol, zafirlukast/budesonide; montelukast/formoterol; montelukast/budesonide; loratadine/montelukast and loratadine/zafirlukast.

Further combinations include tiotropium and fluticasone, tiotropium and budesonide, tiotropium and mometasone, mometasone and salmeterol, formoterol and rofleponide, salmeterol and budesonide, salmeterol and rofleponide, and tiotropium and rofleponide.

5 The above mentioned compounds may exist as salts and esters as well as solvates (hydrates) and all such forms can be included in the formulations of the invention as are racemates and all stereoisomers of the above compounds where such are possible e.g. (R,R)-formoterol and 22R budesonide.

10 The amount of surfactant present in the pharmaceutical aerosol formulation of the present invention may range from about 0.00001% by weight of the total weight of the formulation to about 10% by weight of the total weight of the formulation. Preferably, the amount of surfactant present is at least 0.001% by weight. Preferably, the amount of surfactant present is up to 1% by weight.

15 The formulations of the present invention may contain medicament in an amount of from about 0.01% by weight of the total weight of the formulation to about 10% by weight of the total weight of the formulation. Preferably the medicament is present from about 0.01 to about 1.0 % by weight.

20 The medicament may be dissolved or dispersed in the propellant. If the medicament is to be dispersed in the propellant, preferably the medicament consists largely of particles of 400 nm - 10 μ m, more preferably 400 nm - 6 μ m, especially 500 nm - 5 μ m. In some circumstances particles of at least 0.001 μ m, e.g., at least 0.1 μ m, may be desirable.

25 Preferably at least 50% of the medicament consists of particles within the desired size range. For example at least 60%, preferably at least 70%, more preferably at least 80% and most preferably at least 90% of the medicaments consists of particles within the desired size range.

30 The medicament for use in the present invention may have to be processed prior to inclusion in the formulations, in order to produce particles in the desired size range. For example the medicament may be milled or micronised, for example using suitable equipment, for example an air jet mill, hammer mill, ball mill or using a microfluidiser.

35 Alternatively, particles in the desired particle range may be obtained by for example spray

drying or controlled crystallisation methods, for example crystallisation using supercritical fluids or via an emulsion method, e.g. microfluidisation or homogenisation.

5 In a particular embodiment, the surfaces of the medicament particles may be modified prior to dispersion, for example by spray drying a solution of drug and surfactant or by adsorption of surfactant onto medicament particles. Further techniques for modification of the surfaces of the medicament particles can also be used such as freeze drying, microfluidising and milling.

10 Others ingredients may be added into the formulation of the present invention, if desired. Such ingredients may be for example other pharmaceutically active agents, adjuvants, carriers, flavouring agents, buffers, antioxidants, chemical stabilisers, polymers and other surfactants, for example.

15 The amount of additional ingredients included in the present formulation may be up to 1% w/w.

Dispersion formulations of the present invention may be prepared for example by adding the alkylpolyglycoside surfactant to propellant by pressure- or cold- filling methods,
20 homogenising, and adding medicament. The resultant dispersion may be agitated to ensure dispersion uniformity. Dispersion formulations can also be prepared by co-preparation of particles with excipient, for example by spray drying, to form composite particles.

Solution formulations of the present invention may be prepared by adding the
25 alkylpolyglycoside surfactant and water to propellant by pressure -or cold- filling methods, homogenising, and adding medicament. The mixture is agitated, sonicated or heated until the medicament has dissolved.

Aerosol canisters for use with the formulation of the present invention may comprise, for
30 example, a metal can, for example an aluminium can, closed with a metering valve. In a particular embodiment the internal surfaces of the metal can be coated with, for example, a fluoropolymer. Suitable fluoropolymers include PTFE, PFA, and FEP.

The formulation of the present invention is useful for the local or systemic treatment of
35 diseases and may be administered for example via the upper or lower respiratory tract, including the nasal route. As such the present invention also provides said formulation for

use in therapy; the use of the formulation in the manufacture of a medicament for the treatment of diseases via the respiratory tract; and a method for the treatment of a patient in need of therapy, comprising administering to said patient a therapeutically effective amount of the formulation of the present invention.

5

The invention is illustrated in the following examples.

Formulation preparation

Propellant HFA-134a or HFA-227ea was dispensed chilled (at -55°C) into an open 400ml aluminium can. A 1 inch valve was then crimped onto the can and the propellant allowed to return to ambient temperature (20°C).

Drug (or drug/surfactant mixture) was weighed into a clear polyethyleneterephthalate (PET) vial which was sealed with a non-metering valve. Propellant was then pressure filled through the valve to give the desired drug and surfactant concentrations. The vial was subsequently placed in an ultrasonic bath for one minute to aid dispersion of the particles. 11 samples were prepared according to this method.

Example 1:

Beclomethasone dipropionate (BDP) B.P (SICOR) was weighed (0.2 g) into a 30 ml glass vial and 20 ml of surfactant solution in water (C_{12}G_2 , 0.8 g/l) added. The resultant suspension was incubated in a shaking bath at 25°C for three hours, to allow adsorption of the surfactant to the surface of the drug, and to give a drug-surfactant ratio of 10mg surfactant/g drug. The suspension was centrifuged (15,000 rpm, 20 minutes), and the particles of drug-surfactant were separated from the supernatant and dried in an oven at 50°C for at least 24 hours. The formulation was prepared as described in the preparation paragraph, with the following concentrations:

Beclomethasone dipropionate + C_{12}G_2 : 0.2%

HFA-134a: to 100%

Example 2:

Beclomethasone dipropionate B.P. (SICOR) was weighed (0.2 g) into a 30 ml glass vial and 20 ml of surfactant solution in water (C_{12}G_2 , 0.8 g/l) added. The resultant suspension was incubated in a shaking bath at 25°C for three hours, to allow adsorption of the surfactant to the surface of the drug, and to give a drug-surfactant ratio of 10mg surfactant/g drug. The suspension was then centrifuged (15,000 rpm, 20 minutes) and the particles of drug-surfactant were separated from the supernatant and dried in an oven at 50°C for at least 24 hours. The formulation was prepared as described in the preparation paragraph, with the following concentrations:

Beclomethasone dipropionate + C_2G_2 : 0.2%

HFA-227ea: to 100%

Example 3:

Beclomethasone dipropionate B.P (SICOR) was weighed (0.2 g) into a 30-ml glass vial and 20 ml of surfactant solution in water (C₁₀₋₁₂ d.p. 1.4, 0.8 g/l) added. The resultant suspension was incubated in a shaking bath at 25°C for three hours, to allow adsorption of the surfactant to the surface of the drug, and to give a drug-surfactant ratio of 10 mg surfactant/g drug. The suspension was then centrifuged (15,000 rpm, 20 minutes) and the particles of drug-surfactant were separated from the supernatant and dried in an oven at 50°C for at least 24 hours. The formulation was prepared as described in the preparation paragraph, with the following concentrations:

- 10 Beclomethasone dipropionate + C₁₀₋₁₂ d.p. 1.4: 0.2%
HFA-134a: to 100%

Example 4:

Beclomethasone dipropionate B.P (SICOR) was weighed (0.2 g) into a 30 ml glass vial and 20 ml of surfactant solution in water (C₁₀₋₁₂ d.p. 1.4, 0.8 g/l) added. The resultant suspension was incubated in a shaking bath at 25°C for three hours, to allow adsorption of the surfactant to the surface of the drug, and to give a drug-surfactant ratio of 10 mg surfactant/g drug. The suspension was then centrifuged (15,000 rpm, 20 minutes) and the particles of drug-surfactant were separated from the supernatant and dried in oven at 50°C for at least 24 hours. The formulation was prepared as described in the preparation paragraph, with the following concentrations:

- 20 BDP C₁₀₋₁₂ d.p. 1.4: 0.2%
HFA-227ea: to 100%

Example 5:

Microparticles of salbutamol sulphate-C₁₂₋₁₄ d.p. 1.4 were prepared by spray drying from solution in water using a Büchi 190 mini spray drier fitted with a 7 mm pneumatic nozzle. A 10% w/v salbutamol sulphate in a solution of 0.06 g/l of C₁₂₋₁₄ d.p. 1.4 was spray dried. The conditions and spray drying parameters were: pump speed, 5 ml min⁻¹; air flow rate, 800 l h⁻¹; aspirator level, 5; inlet temperature, 150°C (± 5°C) and outlet temperature 80°C (± 5°C). The material was desiccated immediately after drying. The formulation was prepared as described in the preparation paragraph, with the following concentrations:

- 30 Spray dried (SD) salbutamol sulphate + C₁₂₋₁₄ d.p. 1.4: 0.08%
HFA-134a: to 100%

Example 6:

5 Microparticles of salbutamol sulphate- $C_{12}G_2$ were prepared by spray drying from solution in water using a Büchi 190 mini spray drier fitted with a 7 mm pneumatic nozzle. A 10% w/v salbutamol sulphate in a solution of 0.08 g/l of $C_{12}G_2$ was spray dried. The conditions and spray drying parameters were: pump speed, 5 ml min⁻¹; air flow rate, 800 l h⁻¹; aspirator level, 5; inlet temperature, 150°C (± 5°C) and outlet temperature 80°C (± 5°C). The material was desiccated immediately after drying. The formulation was prepared as described in the preparation paragraph, with the following concentrations:

SD salbutamol sulphate + $C_{12}G_2$: 0.08%

10 HFA-134a: to 100%

Example 7:

Salbutamol sulphate was weighed (0.2 g) into a 30 ml glass vial and 20 ml of surfactant solution in CH_2Cl_2 (C_{14} d.p. 1.4, 0.6 g/l) added. The resultant suspension was incubated in a shaking bath at 25°C for three hours, to allow adsorption of the surfactant to the surface. This was then filtered under vacuum and the particles of drug-surfactant collected and dried overnight at room temperature. The formulation was prepared as described in the preparation paragraph, with the following concentrations:

Salbutamol sulphate + C_{14} d.p. 1.4: 0.08%

20 HFA-134a: to 100%

7

Control 1:

The formulation was prepared as described in the preparation paragraph, with the following concentrations:

25 Beclomethasone dipropionate: 0.2%

HFA-134a: to 100%

Control 2:

The formulation was prepared as described in the preparation paragraph, with the following concentrations:

30 Beclomethasone dipropionate: 0.2%

HFA-227ea: to 100%

Control 3:

35 Microparticles of salbutamol sulphate were prepared by spray drying from an aqueous solution, using a Büchi 190 mini spray drier fitted with a 7 mm pneumatic nozzle. A 10%

w/v salbutamol sulphate solution in water was used. The spray drying parameters were: pump speed 5 ml min⁻¹; air flow rate 800 l h⁻¹; aspirator level 5; inlet temperature 150°C (± 5°C) and outlet temperature 80°C (± 5°C). The material was desiccated immediately after drying. The formulation is then prepared as described in the preparation paragraph, with the following concentrations:

SD salbutamol sulphate: 0.08%

HFA-134a: to 100%

Control 4:

The formulation was prepared as described in the preparation paragraph, with the following concentrations:

Salbutamol sulphate: 0.08%

HFA-134a: to 100%

The following table summarises the list of samples prepared:

Sample	Material	HFA	Surfactant / drug ratio
1	BDP + C ₁₂ G ₂	134a	10mg/1g
2	BDP + C ₁₂ G ₂	227ea	10mg/1g
3	BDP + C ₁₀₋₁₂ d.p. 1.4	134a	10mg/1g
4	BDP + C ₁₀₋₁₂ d.p. 1.4	227ea	10mg/1g
5	SD salbutamol sulphate + C ₁₂₋₁₄ d.p.1.4	134a	0.6mg/1g
6	SD salbutamol sulphate + C ₁₂ G ₂	134a	0.8mg/1g
7	Salbutamol sulphate + C ₁₄ d.p. 1.4	134a	0.8mg/1g
Control 1	BDP	134a	N/A
Control 2	BDP	227ea	N/A
Control 3	SD salbutamol sulphate	134a	N/A
Control 4	Salbutamol sulphate	134a	N/A

Product Evaluation:**OSCAR analysis (Optical suspension characterisation)**

5 In this technique, two matched pairs of infra-red emitter/detector probes are directed at the clear PET vial. The upper detectors were placed 3mm below the suspension surface and the lower detectors placed at 3mm from the bottom of the vial. The infrared probes detect any changes in light transmission through the suspension. Measurement starts automatically when the upper light beam is broken when the vial is inserted. A personal computer then
10 processes the voltage signal and a profile of light transmission is obtained. If the suspension is stable then the transmitted signal will be low. However, if the suspension is unstable and has either settled out or creamed the light transmission will be high. A basic measurement of how rapidly the sample sediments or creams is given by how long it takes the upper (if
15 sedimenting) or lower (if creaming) detector signal to change and reach an equilibrium reading. Further to this, if the fluctuation of the signal with time is great, this is an indication that the sample is flocculating (aggregate size is increasing) and indicates that the suspension is unstable.

Results for the previous samples are listed below. All samples show better stability
20 characteristics than the control samples. Typically, stability times increase from a second timescale to minutes. The presence of the APG does improve substantially the characteristics of the pMDI suspension. The samples also show very good characteristics on storage as demonstrated from the 1 month stability data.

25 Samples 1 and 3 (BDP with APGs in HFA-134a) can be compared with control 1. In both cases the stability of the suspension increases from 10 seconds for the control sample to over one minute for the samples with APGs.

30 Samples 2 and 4 (BDP with APGs in HFA-227) can be compared with control 2. The stability times increase from 20 seconds to well over 10 minutes. In the case of sample 2, the stability is above 60 minutes, indicating an extremely slow creaming rate.

The salbutamol sulphate samples show similar trends, although the timescale for improvements are different. In this case, the best indicators of improvements are the
35 profiles after one month's storage. At the initial timepoint, the suspension stability of salbutamol sulphate formulated with APGs (samples 5 and 6 - SD salbutamol sulphate with

APGs in HFA-134a) are not as good as the control (Control 3) . However after one month's storage, the Control has deteriorated but the suspensions formulated with APGs have improved significantly.

Sample 7 (salbutamol sulphate with APG in HFA-134a) is to be compared with control 7.

- 5 Again, the presence of the APG improves suspension stability by increasing the creaming times, with a particularly beneficial influence after storage.
-




Figure 1 Example 1, BDP+C₁₂G₂ in HFA-134a




Figure 2 Example 2, BDP + C₁₂G₂ in HFA-227ea




Figure 3 Example 3, BDP +C₁₀₋₁₂ d.p. 1.4 in HFA-134a




Figure 4 Example 4, BDP+C₁₀₋₁₂ d.p. 1.4 in HFA-227ea




Figure 5.1 Example 5, SD salbutamol sulphate + C₁₂₋₁₄
d.p. 1.4 in HFA-134a




Figure 5.2 Example 5, SD salbutamol sulphate + C₁₂₋₁₄ d.p. 1.4 in HFA-134a, 1 month stability study




Figure 6.1 Example 6, SD salbutamol sulphate + C₁₂G₂ in HFA-134a




Figure 6.2 Example 6, SD salbutamol sulphate + C₁₂G₂ in HFA-134a, 1 month stability study




Figure 7.1 Example 7, Salbutamol sulphate + C₁₄ d.p. 1.4 in HFA-134a




Figure 7.2 Example 7, Salbutamol sulphate + C₁₄ d.p. 1.4 in HFA-134a, 1 month stability




Figure 8 Control 1, BDP in HFA-134a




Figure 9 Control 2, BDP in HFA-227ea




Figure 10 Control 3, Spray dried salbutamol sulphate in HFA-134a




Figure 11 Control 3, Spray dried salbutamol sulphate in HFA-134a, 1 month stability study




figure 12.1 Control 4, Salbutamol sulphate in HFA-
134a




Figure 12.2 Control 4, Salbutamol sulphate in HFA-134a,
1 month stability

Visual Assessment

- Visual assessment was carried out on the OSCAR samples. After removing the samples from the ultrasonic bath, observations were made to assess the degree of flocculation of the drug suspensions and also the kinetics of sedimentation or creaming.

Sample	Material	HFA	Flocculation ?	Sedimentation /creaming
1	BDP + C ₁₂ G ₂	134a	No	Slow
2	BDP + C ₁₂ G ₂	227ea	No	Slow
3	BDP + C ₁₀₋₁₂ d.p.1.4	134a	No	Very slow
4	BDP + C ₁₀₋₁₂ d.p.1.4	227ea	No	Very slow
5	SD salbutamol sulphate + C ₁₂₋₁₄ d.p. 1.4	134a	Yes	Fast
6	SD salbutamol sulphate + C ₁₂ G ₂	134a	Yes	Fast
7	Salbutamol sulphate + C ₁₄ d.p. 1.4	134a	No	Very slow
Control 1	BDP	134a	Yes	Very fast
Control 2	BDP	227ea	Yes	Very fast
Control 3	SD salbutamol sulphate	134a	Yes	Slow
Control 4	Salbutamol sulphate	134a	Yes	Fast

- For all systems, the presence of APG improved the suspension characteristics. Initial observations are summarised in the previous table. After one month storage, the spray dried salbutamol samples showed a definite improvement compared to the control samples (Control 3).

One month stability:

Sample	Material	HFA	Flocculation ?	Sedimentation /creaming
5	SD salbutamol sulphate + C ₁₂₋₁₄ d.p. 1.4	134a	Yes	slow
6	SD salbutamol sulphate + C ₁₂ G ₂	134a	Yes	Slow
Control 3	SD salbutamol sulphate	134a	Yes	Fast

5 IGC

Surface energy of the drug and the drug surfactant particles was measured by inverse gas chromatography. A silanised U-shaped column (3 mm internal diameter, 30 cm length) was packed with approximately 400 mg of the test powder and dried under nitrogen for 24 hours at 50°C. The powder bed was then allowed to settle for 24 hours at 35°C (temperature used during the experiments) under nitrogen. The test probes used were: hexane, heptane, octane, chloroform, CCL₄, benzene, acetone, tetrahydrofuran, ethyl acetate and diethyl ether. One ml of air containing a minute concentration of each gaseous probe was injected into the column and the retention time measured.

15

The APG reduced the surface energy of the particles notably (not for the SD material?). I think we should remove the SD material & results. The improved suspension characteristics can therefore be attributed to surface modifications.


MATERIAL	γ_s^D MJ/M ² (n=3)	Kd/Ka
BDP	52.34 (1.09)	8.82
BDP + C ₁₂ G ₂	50.25 (1.33)	6.09
BDP + C ₁₀₋₁₂ d.p. 1.4	36.30 (1.62)	2.15
Salbutamol sulphate	43.27 (0.67)	1.76
Salbutamol sulphate + C ₁₄ d.p. 1.4	38.23 (0.36)	2.61

Think we should remove this as the density did change for APG material & the conclusion doesn't match the data.

5

CLAIMS

1. A pharmaceutical aerosol formulation comprising a hydrofluoroalkane propellant, a medicament for inhalation and a surfactant, characterised in that the surfactant is an alkyl-
 5 polyglycoside of formula I:



- wherein DP is the average degree of polymerisation and has a value of from 1 to 4, and R is an alkyl chain or a mixture of alkyl chains having a chain length of from 6 to 22 carbon
 10 atoms; or a derivative thereof.

2. A formulation according to claim 1 in which
 R is 2-ethyl-1-hexylglycoside and DP is 1.6;
 R is a mixture of C₈ and C₁₀ alkyl chains in a ratio of 60 C₈ : 40 C₁₀ and DP is 1.5;
 15 R is a mixture of C₁₆ and C₈ alkyl chains and DP is 1.2 - 1.3; or
 R is a mixture of C₂₀ and C₂₂ alkyl chains and DP is 1.2 - 1.3, or the alkylpolyglycoside is
 n-dodecyl β-D-maltoside (C₁₂G₂), C₁₀ d.p. 2.7, or C₁₀₋₁₂ d.p. 1.4 and C₁₂₋₁₄ d.p.
 20 2.7ea or mixtures thereof.

4. A formulation according to any one of claims 1 to 3 in which the medicament is a β 2-adrenoreceptor agonist; an anticholinergic bronchodilators or a 16, 17-acetal of a pregnane derivative.


5. A formulation according to any one of claims 1 to 3 in which the medicament is formoterol, terbutaline, budesonide or a formoterol/budesonide combination.

6. A formulation according to any one of claims 1 to 3 in which the medicament is a compound of formula (I):



in which

Ar represents a group



A represents a C_{1-12} alkylene chain which may be straight or branched and which is interrupted or terminated by one or more groups selected from -S-, -SO-, SO_2 -, -O-, SO_2NH , NHSO_2 , CR^6R^7 , phenylmethyne, -NH-, -CONH-, -NHCO- and -NHCONH-;

Z represents an aryl group of five or six atoms in a single ring system, which may contain from 1 to 3 heteroatoms selected from N, O and S, which single ring system may be optionally substituted to form a multiple fused ring system of up to 10 atoms, the aryl group being optionally substituted by one or more groups selected from -OH, halogen,

C_{1-6} alkyl, C_{1-6} alkoxy, =O, $-\text{NR}^8\text{R}^9$, or NO_2 ; or a C_{3-12} cycloalkyl group which may contain

from 1 to 3 heteroatoms selected from N, O and S, optionally substituted by one or more groups selected from -OH, halogen, C_{1-6} alkyl, C_{1-6} alkoxy, =O, $-\text{NH}_2$, or NO_2 ;

R^1 , R^2 , R^5 , R^6 , R^7 , R^8 and R^9 each independently represent hydrogen or C_{1-6} alkyl;

and

R^3 and R^4 represent hydrogen, or R^3 and R^4 together form a group -S-, $-\text{NR}^8$ - or $-\text{CH}_2$ -,

and pharmaceutically acceptable derivatives thereof.

7. A formulation according to any one of claims 1 to 3 in which the medicament is

4-Hydroxy-7-[2-[2-[3-(2-phenylethoxy)propylsulphonyl]ethylamino]ethyl]-1,3-benzothiazol-2(3H)-one;

4-Hydroxy-7-[2-[2-[3-(2-phenylethoxy)propoxy]ethylamino]ethyl]-1,3-benzothiazol-2(3H)-one;

5 N-[2-[2-(4-Hydroxy-2-oxo-3H-1,3-benzothiazol-7-yl)ethylamino]ethyl]-2-(phenylethoxy)ethanesulphonamide;

4-Hydroxy-7-[2-[3-[2-[2-(1-naphthalenyl)ethoxy]ethylsulphonyl]propylamino]ethyl]-1,3-benzothiazol-2(3H)-one; and

3-[2-(4-Hydroxy-2-oxo-3H-1,3-benzothiazol-7-yl)ethylamino]-N-[2-[2-(4-methylphenyl)ethoxy]ethyl]propanesulphonamide,
10 and pharmaceutically acceptable salts thereof.

8. A formulation according to any one of claims 1 to 3 in which the medicament is a combination of formoterol/budesonide; formoterol/fluticasone; formoterol/mometasone;
15 salmeterol/fluticasone; formoterol/tiotropium salts; zafirlukast/formoterol, zafirlukast/budesonide; montelukast/formoterol; montelukast/budesonide; loratadine/montelukast and loratadine/zafirlukast, tiotropium and fluticasone, tiotropium and budesonide, tiotropium and mometasone, mometasone and salmeterol, formoterol and rofleponide, salmeterol and budesonide, salmeterol and rofleponide, or tiotropium
20 and rofleponide.

9. A formulation according to any one of claims 1 to 8 in which the amount of surfactant present is at least 0.001% by weight.

25 10. A formulation according to any one of claims 1 to 9 in which the amount of medicament present is from 0.01 to 1.0% by weight.

11. A medicinal aerosol containing a formulation according to any one of claims 1 to 10.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 01/02853

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: A61K 9/12, A61K 47/26

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-INTERNAL, WPI DATA, PAJ, CHEM. ABS DATA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9619198 A1 (ASTRA AKTIEBOLAG), 27 June 1996 (27.06.96) --	1-11
A	STN International, File CAPLUS, CAPLUS accession no. 1997:328726, Document no. 126:306557, Chuo Eazooru Kagaku Kk: "Alkyl glucoside-based aerosol-type foam products"; & JP,A2,09059606, 19970304 --	1-11
A	WO 9830244 A1 (MINNESOTA MINING AND MANUFACTURING COMPANY), 16 July 1998 (16.07.98), page 16, line 17 - page 17, line 7 --	1-11

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

12 March 2002

Date of mailing of the international search report

20-03-2002

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Gerd Strandell/Eö

Telephone No. +46 8 782 25 00

Form PCT/ISA/210 (second sheet) (July 1998)

International application No.
PCT/SE 01/02853

International application No.
PCT/SE 01/02853

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 0019980 A1 (IGEN, INC.), 13 April 2000 (13.04.00), the examples; the claims --	1-11
A	WO 9500151 A1 (UAB RESEARCH FOUNDATION), 5 January 1995 (05.01.95), page 8, line 20 - page 9, line 7; page 11, line 1 - line 5; page 12, line 13 - line 16; page 15, line 19 - line 24; the claims --	1-11
A	WO 9747286 A1 (MINNESOTA MINING AND MANUFACTURING COMPANY), 18 December 1997 (18.12.97), the claims -- -----	1-11

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 01/02853

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	9619198	A1	27/06/96	AU	702880 B	11/03/99
				AU	4359396 A	10/07/96
				BR	9510510 A	07/07/98
				CA	2206782 A	27/06/96
				CN	1170356 A	14/01/98
				CZ	288146 B	16/05/01
				CZ	9701947 A	15/10/97
				EE	9700138 A	15/12/97
				EP	0806940 A	19/11/97
				FI	972655 A	19/06/97
				HU	77775 A	28/08/98
				IL	116460 D	00/00/00
				JP	10510829 T	20/10/98
				NO	972681 A	11/06/97
				NZ	298169 A	29/09/99
				PL	320856 A	10/11/97
				SE	9404469 D	00/00/00
				SK	81197 A	05/11/97
				TR	970136 A	00/00/00
				ZA	9510754 A	24/06/96
				SE	9502452 D	00/00/00
WO	9830244	A1	16/07/98	AU	738444 B	20/09/01
				AU	5709898 A	03/08/98
				EP	0957939 A	24/11/99
				JP	2001508062 T	19/06/01
				US	6019997 A	01/02/00
WO	0019980	A1	13/04/00	EP	1117380 A	25/07/01
				US	6251425 B	26/06/01
WO	9500151	A1	05/01/95	AU	7177794 A	17/01/95
				US	5661130 A	26/08/97
WO	9747286	A1	18/12/97	AU	726382 B	02/11/00
				AU	3373997 A	07/01/98
				CA	2257841 A	18/12/97
				EP	0934057 A	11/08/99
				GB	9612297 D	00/00/00
				JP	2000513340 T	10/10/00
				NO	985720 A	11/02/99
				US	6054488 A	25/04/00
				ZA	9704546 A	23/11/98

Form PCT/ISA/210 (patent family annex) (July 1998)